

2011 Cow Herd Protocol

Pfizer Animal Health

Yearling Heifers:

- ❖ Pre-breeding (30 – 60 days: 4 weeks minimum prior to breeding)
 - Vaccines
 - Bovi-shield Gold FP5 VL5 HB (5 ml SQ)
 - Ultrabac 7 (5 ml SQ)
 - Internal Parasite Control
 - Dectomax injectable (1 ml/110 lb SQ) or Dectomax Pour-On

Bred Heifers:

- ❖ Pregnancy Exam
 - Vaccines
 - Leptoferm-5 (2 ml IM)
 - Internal Parasite Control
 - Dectomax injectable (1 ml/110 lb SQ) or Dectomax Pour-On

Optional:

Scours Control (Cows and Bred Heifers Pre-Calving):

- ❖ Pre-Calving vaccination (3 to 6 weeks prior to calving)
 - ScourGuard 4KC (2 ml IM)
 - Note: For first calf heifers and any cows not previously vaccinated with ScourGuard 4KC, the scour vaccine must be administered twice approximately 3 weeks apart with an initial vaccination administered 3 weeks prior to this vaccination

(Over)

Mature Cow Herd:

- ❖ Pre-breeding (3– 4 weeks)
 - Vaccines
 - Bovi-Shield Gold FP5 VL5 HB (5 ml SQ)*
 - Internal Parasite Control
 - Dectomax injectable (1 ml/110 lb SQ) or Dectomax Pour-On

- ❖ Pregnancy Exam (1 – 4 months gestation)
 - Vaccines
 - Leptoferm 5 (2 ml IM)
 - Internal Parasite Control
 - Dectomax Injectable or Dectomax Pour-On (seasonal & timing consideration)

Jennie Schutte
646.306.6131
Jennie.schutte@pfizer.com

Please consult with your veterinarian on this protocol.

(Over)