



2016 Cow Herd Protocol Zoetis Animal Health

Yearling Heifers:

Pre-breeding (30 – 60 days: 4 weeks minimum prior to breeding)

-Vaccines

- Bovi-shield Gold FP5 VL5 HB (5 ml SQ)*
- Ultrabac 7 (5 ml SQ)

-Internal Parasite Control

- Dectomax injectable (1 ml/110 lb SQ) or Dectomax Pour-On

Bred Heifers:

Pregnancy Exam

-Vaccines

- Leptoferm-5 (2 ml IM)

- Internal Parasite Control

- Dectomax injectable (1 ml/110 lb SQ) or Dectomax Pour-On

Optional:

Scours Control (Cows and Bred Heifers Pre-Calving):

-Pre-Calving vaccination (3 to 6 weeks prior to calving)

- ScourGuard 4KC (2 ml IM)
- Note: For first calf heifers and any cows not previously vaccinated with ScourGuard 4KC, the scour vaccine must be administered twice approximately 3 weeks apart with an initial vaccination administered 6-9 weeks prior to calving and the booster dose 3-6 weeks prior to calving.

(Over)

Mature Cow Herd:

Pre-breeding (3– 4 weeks)

-Vaccines

- Bovi-Shield Gold FP5 VL5 HB (5 ml SQ)*

-Internal Parasite Control

- Dectomax injectable (1 ml/110 lb SQ) or Dectomax Pour-On

Pregnancy Exam (1 – 4 months gestation)

-Vaccines

- Leptoferm 5 (2 ml IM)

-Internal Parasite Control

- Dectomax Injectable or Dectomax Pour-On (seasonal & timing consideration)

Josh Wackler
541.740.7475
Josh.wackler@zoetis.com

***Please consult with your veterinarian on this protocol**